

N-[3,4-dimethoxycinnamoyl]-anthranilic acid (tranilast) suppresses microglial inducible nitric oxide synthase (iNOS) expression and activity induced by interferon- γ (IFN- γ)

*¹Michael Platten, ¹Wolfgang Wick, ¹Jörg Wischhusen & ¹Michael Weller

¹Laboratory of Molecular Neuro-Oncology, Department of Neurology, University of Tübingen, School of Medicine, Hoppe-Seyler Str. 3, 72076 Tübingen, Germany

1 Microglial cells up-regulate inducible nitric oxide synthase (iNOS) expression in response to various pro-inflammatory stimuli including interferon- γ (IFN- γ), allowing for the release of nitric oxide (NO). Tranilast (N-[3,4-dimethoxycinnamoyl]-anthranilic acid) is an antiallergic compound with suppressive effects on the activation of monocytes.

2 Here, we show that N9 murine microglial cells express iNOS mRNA and protein and release nitric oxide into the culture medium in response to IFN- γ (200 u ml⁻¹) as measured by Northern and Western blot analyses and Griess assay.

3 Exposure to non-toxic doses of tranilast (30–300 μ M) leads to a concentration-dependent inhibition of IFN- γ -induced (200 u ml⁻¹) iNOS mRNA and protein expression. This is paralleled by a suppression of NO-release into the cell culture medium.

4 Inhibition of IFN- γ -induced iNOS mRNA expression by tranilast is paralleled by an inhibition of nuclear factor- κ B (NF- κ B) activation and phosphorylation of inhibitory κ B (I κ B) as determined by Western blot analyses and NF- κ B reporter gene assay.

5 These results suggest that tranilast-mediated suppression of microglial iNOS activity induced by IFN- γ involves the inhibition of NF- κ B-dependent iNOS mRNA expression.

British Journal of Pharmacology (2001) **134**, 1279–1284

Keywords: Tranilast; microglia; nitric oxide; inducible nitric oxide synthase; interferon- γ ; nuclear factor- κ B; inhibitory- κ B

Abbreviations: ALS, amyotrophic lateral sclerosis; CNS, central nervous system; COX-2, cyclo-oxygenase type 2; DEX, dexamethasone; IFN- γ , interferon- γ ; IL-1 β , interleukin-1 β ; I κ B, inhibitory κ B; iNOS, inducible nitric oxide synthase; LPS, lipopolysaccharide; NF- κ B, nuclear factor κ B; NO, nitric oxide; PGE₂, prostaglandin E₂; TGF- β , transforming growth factor- β ; TNF- α , tumour necrosis factor- α ; TNL, tranilast

Introduction

Microglial cells share many morphological and functional features with monocytes and macrophages and are thought to contribute to inflammatory processes in the brain. Microglia is activated during brain ischaemia, infection, demyelination, neurodegeneration and tumorigenesis (Gonzalez-Scarano & Baltuch, 1999). On a cellular level, activation of microglial cells, like monocytic cells, results in the expression of pro-inflammatory mediators such as cyclo-oxygenase type 2 (COX-2), tumour necrosis factor- α (TNF- α) and nitric oxide (NO). NO is a free radical generated from L-arginine by NO synthases (NOS) (Griffith & Stuehr, 1995). It is involved in a wide range of physiological and pathological processes including the regulation of vascular homeostasis, neurotransmission and inflammation (Nathan, 1992). Three isoforms of NOS have been characterized. Neuronal (type I or nNOS) and endothelial (type III or eNOS) NOS are constitutively expressed and are mainly regulated posttranscriptionally by calmodulin-dependent pathways. The inducible NOS isoform (type II or iNOS) is not expressed in healthy tissues but is rapidly expressed *de novo* in many cell types including

astrocytes and microglia in response to lipopolysaccharide (LPS), a bacterial wall component, or inflammatory cytokines such as interleukin-1 β (IL-1 β), TNF- α or IFN- γ (Galea *et al.*, 1992; Hu *et al.*, 1995; Nathan & Xie, 1994). When produced in excess amounts NO may promote neuronal death by reacting with superoxide anion to generate peroxynitrite (Beckman *et al.*, 1990). Consequently, overproduction of NO has been implicated in neurological disorders associated with neuronal damage including brain ischaemia, Parkinson's disease, Alzheimer's disease and amyotrophic lateral sclerosis (ALS) (Chabrier *et al.*, 1999). Accordingly, pharmacological inhibition of iNOS has been shown to attenuate brain damage in an animal model of brain ischaemia (Nagayama *et al.*, 1998).

Tranilast (N-[3,4-dimethoxycinnamoyl] anthranilic acid) is an orally active anti-allergic compound which is clinically effective in the control of autoimmune diseases such as bronchial asthma and atopic dermatitis, presumably by inhibiting the release of various chemical mediators from mast cells (Azuma *et al.*, 1976). In addition, tranilast has been shown to prevent restenosis after percutaneous transluminal angioplasty in patients with coronary heart disease (Tamai *et al.*, 1999). Recently, our laboratory has defined a role for tranilast as a novel anti-glioma agent (Platten *et al.*, 2001). The

*Author for correspondence;
E-mail: michael.platten@uni-tuebingen.de

molecular effects of tranilast are not completely understood. The inhibitory effects of tranilast on collagen accumulation, proliferation and migration of different cell types may be attributed to antagonizing TGF- β (Fukuyama *et al.*, 1996; Miyazawa *et al.*, 1995). The suppressive effects on fibrosis may involve the inhibition of activated macrophages known to release NO (Kawano & Noma, 1993; Mori *et al.*, 1995; Suzawa *et al.*, 1992). On the other hand, there are lines of evidence that tranilast may be a positive regulator of iNOS expression and activity in vascular smooth muscle cells and rat mesangial cells (Hishikawa *et al.*, 1996).

Here we investigate the effects of tranilast on iNOS expression and activity in microglial cells.

Methods

Reagents and cell culture

Tranilast was a generous gift of Kissei Pharmaceuticals (Nagano, Japan). Murine recombinant IFN- γ was purchased from Boehringer (Mannheim, Germany). Dexamethasone (DEX) was obtained from Sigma (Deisenhofen, Germany). N9 murine microglial cells were a kind gift from P. Ricciardi-Castagnoli (Milano, Italy) (Righi *et al.*, 1989). The cells were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% foetal calf serum (FCS), 2 mM glutamine and penicillin (100 u ml⁻¹)/streptomycin (100 μ g ml⁻¹), and maintained at 37°C in an atmosphere of 5% CO₂. Viability was assessed by crystal violet staining.

iNOS activity

iNOS activity was assessed by the Griess assay as described elsewhere (Rieger *et al.*, 1998). Briefly, conditioned supernatant (100 μ l) was incubated with an equal volume of 1% sulphanilamide/0.1% naphthyl ethylene diamine dihydrochloride/2.5% H₃PO₄ (Griess reagent) (Green *et al.*, 1982), incubated for 5 min at room temperature, and the absorbance at 546 nm was determined. Solutions of NaNO₂ diluted in DMEM served as standards. The absorbance readings resulting from DMEM alone were subtracted from the sample values. Cell density was assessed using crystal violet staining to control for differences in cell proliferation. iNOS activity is thus expressed here as NO₂ accumulated per 48 h per 10⁵ cells.

Immunoblot analysis

For the preparation of whole cell lysates, the cells were rinsed in phosphate-buffered saline (PBS), harvested, centrifuged at 1200 $\times g$, lysed in 0.1 M TRIS-HCl (pH 7.2) containing 0.1% NP40, 0.1 mM EDTA and 5 μ g ml⁻¹ PMSF for 40 min on ice, and centrifuged at 10,000 $\times g$ for 10 min. Protein concentration was determined using Biorad reagents with photometric analysis. Proteins (20 μ g lane⁻¹) were separated by 10–12% SDS-PAGE and electroblotted on nitrocellulose. Equal protein loading was controlled by Ponceau S staining. After blocking for 1 h in PBS supplemented with 5% skimmed milk and 0.1% Tween 20, immunodetection was performed using anti-iNOS rabbit polyclonal antibody (1:1000, Cayman, Ann Arbor, MI, U.S.A.) and anti-

phospho-I κ B α mouse monoclonal antibody (1:1000, New England Biolabs, Frankfurt am Main, Germany). Bands were visualized using horseradish peroxidase (HRP)-conjugated anti-rabbit IgG (1:4000, Santa Cruz Biotechnology) or anti-mouse IgG (1:4000, Amersham) and enhanced chemoluminescence (ECL) (Amersham).

Northern blot analysis

Total RNA was isolated using the RNeasy kit (Qiagen, Hilden, Germany). Total RNA (10 μ g) was separated on 1.2% agarose gels and blotted onto nylon membranes (Amersham). The filters were hybridized according to standard procedures (Platten *et al.*, 2001) with a ³²P-labelled murine cDNA probe for iNOS. For the generation of the iNOS probe, N9 cells were stimulated with IFN- γ (200 u ml⁻¹) for 24 h. Total RNA (5 μ g) was subjected to reverse transcription using SuperScript II (Gibco-BRL, Gaithersburg, MD, U.S.A.) and oligo-dT priming (Amersham Pharmacia Biotech, Uppsala, Sweden). iNOS fragments were PCR-amplified using primers 5'-AAGCTGCATGTGACATCGAC-3' and 5'-ATGTGTCTGCAGATGTGCTG-3' corresponding to nucleotides 386–405 and 839–858 of murine iNOS cDNA. Equal loading was assured by ethidium bromide staining.

NF- κ B reporter assay

N9 murine microglial cells were seeded in a 96-well plate (8 \times 10³ well⁻¹) and allowed to adhere for 24 h. The cells were transfected with an NF- κ B cis-reporter gene plasmid (PathDetect[®] no. 219077, Stratagene) which encodes firefly luciferase in a NF- κ B-dependent-manner. For each transfection, 0.2 μ g DNA and 0.6 μ l FuGene (Roche, Mannheim, Germany) were used. The cells were either co-transfected with 0.02 μ g of the pFC-Mekk positive control plasmid, included in the NF- κ B cis-reporting system from Stratagene, or with the pcDNA-3 plasmid, to maintain a constant amount of total DNA. In order to assess the unspecific background, a control transfection with pcDNA-3 only was included in all experiments. After 24 h the cells were washed twice with PBS (0.12 M NaCl, 0.01 M NaH₂PO₄ \times H₂O, 0.031 M K₂HPO₄) and lysed using 40 μ l well⁻¹ of Cell Lysis Buffer (mM): tricine pH 7.8 40, NaCl 50, EDTA 2, MgSO₄ 1, DTT 5, 1% Triton[®] X-100, all from Sigma. To optimize lysis, a freeze-thaw cycle was performed. Then the lysate was transferred to a LumiNunc TM plate (Nunc, Roskilde, Denmark). Luciferase assay reagent (100 μ l) containing (mM): tricine pH 7.8 40, MgSO₄ 10, EDTA 0.5, DTT 10, coenzyme A 0.5 and beetle luciferin 0.5, (0.5 M ATP), (all from Sigma) were added and luminescence was measured in a LumimatPlus (EG&G Berthold, Pforzheim, Germany). Background was subtracted from all values and the remaining luciferase activity was expressed in per cent of the positive control (pFC-Mekk transfection). Thus, the results become independent from transfection efficiency. The value of the positive control remained largely unchanged through all treatment groups.

Statistical analyses

Experiments were usually performed in triplicate and repeated three times. The significance was evaluated by *t*-test or ANOVA at *P* < 0.05, *P* < 0.01 or *P* < 0.001.

Results

Tranilast suppresses microglial iNOS protein expression and activity induced by IFN- γ

While N9 microglial cells displayed no constitutive iNOS protein expression, incubation with IFN- γ (200 u ml⁻¹) induced strong iNOS protein expression in N9 cells in a time-dependent manner as measured by immunoblot analysis. iNOS protein was weakly detectable 4 h after stimulation, strongly induced after 8 h and lasted for at least 24 h (Figure 1A). Addition of tranilast (300 μ M) greatly suppressed IFN- γ -induced iNOS protein expression as did dexamethasone (500 nM) (Figure 1B). No detectable amounts of NO₂ were measured in the conditioned medium of untreated N9 cells indicating that there was no constitutive iNOS activity (data not shown). When stimulated with IFN- γ (200 u ml⁻¹) the cells released 13 μ M (\pm 3 μ M) NO₂ per 48 h per 10⁵ cells. Addition of tranilast led to a concentration-dependent inhibition of NO release with a reduction by 65% at 300 μ M. In comparison, addition of 500 nM dexamethasone reduced the amount of NO₂ by 75% (Figure 1C).

Tranilast suppresses microglial iNOS mRNA expression induced by IFN- γ

IFN- γ (200 u ml⁻¹) induced strong iNOS mRNA expression in N9 cells in a time-dependent manner as assessed by Northern Blot (Figure 2A). Coincubation with tranilast (300 μ M) greatly suppressed iNOS mRNA expression, indicating that tranilast inhibits iNOS induction on a transcriptional level (Figure 2B).

Tranilast suppresses NF- κ B activation induced by IFN- γ in N9 microglia

Since NF- κ B is an essential transcription factor for the induction of iNOS gene expression, we next asked whether tranilast modulates the activation of NF- κ B in response to IFN- γ . At 24 h after transfection with an NF- κ B reporter gene plasmid, incubation with IFN- γ (200 u ml⁻¹) led to a time-dependent activation of luciferase activity. Of note, there was a small but significant increase of luciferase activity 4 h after stimulation with IFN- γ which peaked after 30 h. There was an induction of protein expression of phosphorylated I κ B by IFN- γ preceding the activation of NF- κ B reporter gene activity as measured by immunoblot analysis (Figure 3A). When coincubated with tranilast, there was a concentration-dependent suppression of I κ B phosphorylation and NF- κ B reporter gene activity (Figure 3B).

Discussion

De novo expression of iNOS and the release of large amounts of NO by microglia is considered to play a significant role in the pathogenesis of various neurological disorders. For instance, induction of microglial NO synthesis is thought to contribute to the neuronal damage in AIDS dementia (Adamson *et al.*, 1996), Alzheimer's disease (Goodwin *et al.*, 1995) and Parkinson's disease (Dehmer *et al.*, 2000). Further, microglial-derived NO is thought to play a crucial




Figure 1 Tranilast suppresses iNOS protein expression and activity in N9 microglia in response to IFN- γ . (A) N9 cells were incubated with IFN- γ (200 u ml⁻¹) and protein lysates were prepared at the time points indicated. Immunoblot analysis was performed using an anti-iNOS antibody revealing a band at ~120 kDa. (B) N9 cells were incubated with medium alone, IFN- γ alone (200 u ml⁻¹) or coincubated with IFN- γ and tranilast (TNL, 300 μ M) or dexamethasone (DEX, 500 nM) for 24 h. (C) N9 cells were stimulated with IFN- γ alone or in combination with dexamethasone (DEX, upper panel) or tranilast (TNL, lower panel) at the concentrations indicated. Supernatant was collected after 48 h and nitrite was measured using the Griess assay. Values are expressed as NO₂ accumulated per 10⁵ cells (mean and s.e.mean, n = 3, * P < 0.05, ** P < 0.01, *** P < 0.001).

role in oligodendrocyte cytotoxicity and demyelination in multiple sclerosis (Merrill *et al.*, 1993). Inhibition of iNOS activity in microglial cells may thus represent an interesting target to ameliorate cytotoxic effects in various diseases associated with increased NO production.

The present study seeks to investigate the effects of tranilast, an orally active anti-allergic drug, on the expression and activity of iNOS in murine N9 microglial cells. In macrophages, high doses of IFN- γ initiate the synthesis of NO through the engagement of response elements in the promoter region of the iNOS gene. Induction of iNOS in murine macrophages by IFN- γ alone appears to be strictly dependent on the activation of the transcription factor NF- κ B (Lowenstein *et al.*, 1993). Activation of NF- κ B requires




Figure 2 Tranilast suppresses iNOS mRNA expression in N9 microglia in response to IFN- γ . (A) N9 cells were incubated with IFN- γ (200 u ml $^{-1}$) and mRNA was prepared at the time points indicated. Northern blot analysis was performed using an iNOS cDNA probe revealing a band running at \sim 1.2 kB. Equal loading was assured by ethidium bromide staining of the 18 S and 28 S ribosomal fractions. (B) N9 cells were incubated with tranilast (30–300 μ M) and stimulated with IFN- γ (200 u ml $^{-1}$) for 24 h. Northern blot analysis was performed as in (A).




Figure 3 Tranilast suppresses NF- κ B activation and I κ B α phosphorylation in N9 microglia in response to IFN- γ . N9 cells were incubated with IFN- γ (200 u ml $^{-1}$) with or without a 4 h preincubation with tranilast (30–300 μ M) and protein lysates were prepared at the time points indicated. Immunoblot analysis was performed using an anti-phospho-I κ B α antibody (A, B, upper panels). In parallel, N9 cells were transiently transfected with an NF- κ B reporter plasmid or MEKK plasmid serving as a positive control. After 24 h cells were incubated with IFN- γ (200 u ml $^{-1}$) with or without a 4 h preincubation with tranilast (30–300 μ M). Luciferase activity was determined as described. Values are expressed as per cent relative light units of the positive control. Data are expressed as mean and s.e.mean ($n=3$; * $P<0.05$, ** $P<0.01$, *** $P<0.001$, effect of IFN- γ compared to control (A); * $P<0.05$, ** $P<0.01$, effect of IFN- γ + TNL compared with IFN- γ alone (B)) (A, B, lower panel).

phosphorylation on specific serine residues and degradation of I κ B α which then dissociates from the NF- κ B/I κ B complex and allows NF- κ B to translocate to the nucleus and bind to κ B motifs (Mercurio *et al.*, 1997; Woronicz *et al.*, 1997).

We find, that incubation of N9 microglial cells with IFN- γ (200 u ml⁻¹) leads to an induction of iNOS mRNA and protein expression accompanied by a release of NO into the cell culture medium (Figures 1A and 2A). After transient transfection with an NF- κ B reporter gene plasmid, there was a time-dependent induction of luciferase activity upon stimulation with IFN- γ indicating an activation of transcriptional activity of NF- κ B. Activation of NF- κ B by IFN- γ appears to be achieved by degradation of I κ B since IFN- γ induces phosphorylation of I κ B α in N9 microglia (Figure 3A).

We further show that the anti-allergic drug tranilast inhibits the expression of iNOS (Figures 1B and 2B) and the release of NO from N9 microglia (Figure 1C). Since NF- κ B activation is involved in iNOS induction we tested the hypothesis that tranilast blocks NF- κ B activation by transiently transfecting N9 cells with an NF- κ B reporter plasmid. Our data suggest that the inhibition of iNOS

mRNA expression involves suppression of NF- κ B activation (Figure 3B). This is supported by several studies demonstrating inhibitory effects of tranilast on the transcription of other genes transcriptionally activated by NF- κ B binding, such as IL-5 and COX-2 (Hiratochi *et al.*, 2000; Inoue *et al.*, 1997).

How tranilast blocks IFN- γ -induced NF- κ B activation, however, is not clear. Like most inhibitors of NF- κ B activation, tranilast prevents IFN- γ -induced phosphorylation and degradation of I κ B α (Figure 3B). The signalling events involved in phosphorylation of I κ B include multiple protein kinases belonging to the family of MAPK (Karin & Delhase, 1998), PKC and PKB (Bauer *et al.*, 2001), all of which may be suppressed by tranilast (Koyama *et al.*, 1999; Watanabe *et al.*, 2000). The exact mechanisms leading to the suppression of NF- κ B activation by tranilast require further investigation.

Taken together, we demonstrate that tranilast suppresses the induction of iNOS expression and activity by IFN- γ in N9 microglia, possibly as a result of preventing NF- κ B activation, and thus identify tranilast as a novel therapeutic agent for various neurological conditions associated with increased NO production, including multiple sclerosis, cerebral ischaemia, Alzheimer's disease and AIDS dementia.

References

- ADAMSON, D.C., WILDEMANN, B., SASAKI, M., GLASS, J.D., MCARTHUR, J.C., CHRISTOV, V.I., DAWSON, T.M. & DAWSON, V.L. (1996). Immunologic NO synthase: elevation in severe AIDS dementia and induction by HIV-1 gp41. *Science*, **274**, 1917–1921.
- AZUMA, H., BANNO, K. & YOSHIMURA, T. (1976). Pharmacological properties of N-(3',4'-dimethoxycinnamoyl) anthranilic acid (N-5'), a new anti-atopic agent. *Br. J. Pharmacol.*, **58**, 483–488.
- BAUER, B., KRUMBOCK, N., FRESSER, F., HOCHHOLDINGER, F., SPITALER, M., SIMM, A., UBERALL, F., SCHRAVEN, B. & BAIER, G. (2001). Complex formation and cooperation of PKC {theta} and Akt1/PKB {alpha} in the NF- κ B transactivation cascade in Jurkat T cells. *J. Biol. Chem.*, **276**, 31627–31634.
- BECKMAN, J.S., BECKMAN, T.W., CHEN, J., MARSHALL, P.A. & FREEMAN, B.A. (1990). Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc. Natl. Acad. Sci. U.S.A.*, **87**, 1620–1624.
- CHABRIER, P.E., DEMERLE-PALLARDY, C. & AUGUET, M. (1999). Nitric oxide synthases: targets for therapeutic strategies in neurological diseases. *Cell. Mol. Life Sci.*, **55**, 1029–1035.
- DEHMER, T., LINDENAU, J., HAID, S., DICHGANS, J. & SCHULZ, J.B. (2000). Deficiency of inducible nitric oxide synthase protects against MPTP toxicity in vivo. *J. Neurochem.*, **74**, 2213–2216.
- FUKUYAMA, J., MIYAZAWA, K., HAMANO, S. & UJIE, A. (1996). Inhibitory effects of tranilast on proliferation, migration, and collagen synthesis of human vascular smooth muscle cells. *Can. J. Physiol. Pharmacol.*, **74**, 80–84.
- GALEA, E., FEINSTEIN, D.L. & REIS, D.J. (1992). Induction of calcium-independent nitric oxide synthase activity in primary rat glial cultures. *Proc. Natl. Acad. Sci. U.S.A.*, **89**, 10945–10949.
- GONZALEZ-SCARANO, F. & BALTUCH, G. (1999). Microglia as mediators of inflammatory and degenerative diseases. *Annu. Rev. Neurosci.*, **22**, 219–240.
- GOODWIN, J.L., UEMURA, E. & CUNNICK, J.E. (1995). Microglial release of nitric oxide by the synergistic action of beta-amyloid and IFN-gamma. *Brain Res.*, **692**, 207–214.
- GREEN, L.C., WAGNER, D.A., GLOGOWSKI, J., SKIPPER, P.L., WISHNOK, J.S. & TANNENBAUM, S.R. (1982). Analysis of nitrate, nitrite, and [15N]nitrate in biological fluids. *Anal. Biochem.*, **126**, 131–138.
- GRIFFITH, O.W. & STUEHR, D.J. (1995). Nitric oxide synthases: properties and catalytic mechanism. *Annu. Rev. Physiol.*, **57**, 707–736.
- HIRATOCHI, M., TAKAMOTO, M., TATEMACHI, S. & SUGANE, K. (2000). Inhibition of interleukin 5 production with no influence on interleukin 4 production by an anti-allergic drug, tranilast, in *Toxocara canis*-infected mice. *Int. J. Immunopharmacol.*, **22**, 463–471.
- HISHIKAWA, K., NAKAKI, T., HIRAHASHI, J., MARUMO, T. & SARUTA, T. (1996). Tranilast restores cytokine-induced nitric oxide production against platelet-derived growth factor in vascular smooth muscle cells. *J. Cardiovasc. Pharmacol.*, **28**, 200–207.
- HU, S., SHENG, W.S., PETERSON, P.K. & CHAO, C.C. (1995). Differential regulation by cytokines of human astrocyte nitric oxide production. *Glia*, **15**, 491–494.
- INOUE, H., OHSHIMA, H., KONO, H., YAMANAKA, M., KUBOTA, T., AIHARA, M., HIROI, T., YAGO, N. & ISHIDA, H. (1997). Suppressive effects of tranilast on the expression of inducible cyclooxygenase (COX2) in interleukin-1beta-stimulated fibroblasts. *Biochem. Pharmacol.*, **53**, 1941–1944.
- KARIN, M. & DELHASE, M. (1998). JNK or IKK, AP-1 or NF- κ B, which are the targets for MEK kinase 1 action? *Proc. Natl. Acad. Sci. U.S.A.*, **95**, 9067–9069.
- KAWANO, Y. & NOMA, T. (1993). Cell action mechanism of tranilast—effect on the expression of HLA-class II antigen. *Int. J. Immunopharmacol.*, **15**, 487–500.
- KOYAMA, S., TAKAGI, H., OTANI, A., SUZUMA, K., NISHIMURA, K. & HONDA, Y. (1999). Tranilast inhibits protein kinase C-dependent signalling pathway linked to angiogenic activities and gene expression of retinal microcapillary endothelial cells. *Br. J. Pharmacol.*, **127**, 537–545.
- LOWENSTEIN, C.J., ALLEY, E.W., RAVAL, P., SNOWMAN, A.M., SNYDER, S.H., RUSSELL, S.W. & MURPHY, W.J. (1993). Macrophage nitric oxide synthase gene: two upstream regions mediate induction by interferon gamma and lipopolysaccharide. *Proc. Natl. Acad. Sci. U.S.A.*, **90**, 9730–9734.
- MERCURIO, F., ZHU, H., MURRAY, B.W., SHEVCHENKO, A., BENNETT, B.L., LI, J., YOUNG, D.B., BARBOSA, M., MANN, M., MANNING, A. & RAO, A. (1997). IKK-1 and IKK-2: cytokine-activated I κ B kinases essential for NF- κ B activation. *Science*, **278**, 860–866.
- MERRILL, J.E., IGNARRO, L.J., SHERMAN, M.P., MELINEK, J. & LANE, T.E. (1993). Microglial cell cytotoxicity of oligodendrocytes is mediated through nitric oxide. *J. Immunol.*, **151**, 2132–2141.

- MIYAZAWA, K., KIKUCHI, S., FUKUYAMA, J., HAMANO, S. & UJIIE, A. (1995). Inhibition of PDGF- and TGF-beta 1-induced collagen synthesis, migration and proliferation by tranilast in vascular smooth muscle cells from spontaneously hypertensive rats. *Atherosclerosis*, **118**, 213–221.
- MORI, H., TANAKA, H., KAWADA, K., NAGAI, H. & KODA, A. (1995). Suppressive effects of tranilast on pulmonary fibrosis and activation of alveolar macrophages in mice treated with bleomycin: role of alveolar macrophages in the fibrosis. *Jpn. J. Pharmacol.*, **67**, 279–289.
- NAGAYAMA, M., ZHANG, F. & IADECOLA, C. (1998). Delayed treatment with aminoguanidine decreases focal cerebral ischemic damage and enhances neurologic recovery in rats. *J. Cereb. Blood Flow. Metab.*, **18**, 1107–1113.
- NATHAN, C. (1992). Nitric oxide as a secretory product of mammalian cells. *FASEB. J.*, **6**, 3051–3064.
- NATHAN, C. & XIE, Q.W. (1994). Regulation of biosynthesis of nitric oxide. *J. Biol. Chem.*, **269**, 13725–13728.
- PLATTEN, M., WILD-BODE, C., WICK, W., LEITLEIN, J., DICHGANS, J. & WELLER, M. (2001). N-[3,4-dimethoxycinnamoyl]-anthranilic acid (tranilast) inhibits transforming growth factor-beta release and reduces migration and invasiveness of human malignant glioma cells. *Int. J. Cancer*, **93**, 53–61.
- RIEGER, J., STANDER, M., LOSCHMANN, P.A., HENEKA, M., DICHGANS, J., KLOCKGETHER, T. & WELLER, M. (1998). Synthesis and biological effects of NO in malignant glioma cells: modulation by cytokines including CD95L and TGF-beta, dexamethasone, and p53 gene transfer. *Oncogene*, **17**, 2323–2332.
- RIGHI, M., MORI, L., DE LIBERO, G., SIRONI, M., BIONDI, A., MANTOVANI, A., DONINI, S.D. & RICCIARDI-CASTAGNOLI, P. (1989). Monokine production by microglial cell clones. *Eur. J. Immunol.*, **19**, 1443–1448.
- SUZAWA, H., KIKUCHI, S., ICHIKAWA, K. & KODA, A. (1992). Inhibitory action of tranilast, an anti-allergic drug, on the release of cytokines and PGE2 from human monocytes-macrophages. *Jpn. J. Pharmacol.*, **60**, 85–90.
- TAMAI, H., KATOH, O., SUZUKI, S., FUJII, K., AIZAWA, T., TAKASE, S., KUROGANE, H., NISHIKAWA, H., SONE, T., SAKAI, K. & SUZUKI, T. (1999). Impact of tranilast on restenosis after coronary angioplasty: tranilast restenosis following angioplasty trial (TREAT). *Am. Heart. J.*, **138**, 968–975.
- WATANABE, S., MATSUDA, A., SUZUKI, Y., KONDO, K., IKEDA, Y., HASHIMOTO, H. & UMEMURA, K. (2000). Inhibitory mechanism of tranilast in human coronary artery smooth muscle cells proliferation, due to blockade of PDGF-BB-receptors. *Br. J. Pharmacol.*, **130**, 307–314.
- WORONICZ, J.D., GAO, X., CAO, Z., ROTHE, M. & GOEDEL, D.V. (1997). IkappaB kinase-beta: NF-kappaB activation and complex formation with IkappaB kinase-alpha and NIK. *Science*, **278**, 866–869.

(Received August 8, 2001
Accepted September 5, 2001)